

Lactating Intramammary Treating Options				Indications						Rx Required	
Product	Mfr.	Milk Withhold Post Calving	Slaughter Withhold	Coagulase-neg. staph	Staph aureus	Strep agalactiae	Strep dysgalactiae	Strep uberis	E. coli		Dosage
<b>Amoxi-Mast®</b> <i>(amoxicillin)</i>	Merck	60 Hours	12 Days		X	X				1 syringe; may repeat every 12 hrs for a maximum of 3 treatments	X
<b>Dariclox®</b> <i>(sodium cloxacillin)</i>	Merck	48 Hours	10 Days		X	X				1 syringe; may repeat every 12 hrs for a maximum of 3 treatments	X
<b>Masti-Clear®</b> <i>(penicillin)</i>	Hanford's US Vet	60 Hours	3 Days			X	X	X		1 syringe; may repeat every 12 hrs for a maximum of 3 treatments	
<b>PolyMast™</b> <i>(hetacillin potassium)</i>	B.I.V.I.	72 Hours	10 Days		X	X	X		X	1 syringe every 24 hr for a maximum of 3 treatments	X
<b>SPECTRAMAST® LC</b> <i>(ceftiofur hydrochloride)</i>	zoetis™	72 Hours	2 Days	X			X		X	1 syringe every 24 hrs; may repeat for up to 8 consecutive days	X
<b>ToDAY®</b> <i>(cephapirin sodium)</i>	B.I.V.I.	96 Hours	4 Days		X	X				1 syringe every 12 hrs for a maximum of 2 treatments	

\*9-day slaughter withhold required after last treatment when used for 2 days.  
21-day slaughter withhold when treatment period has been 3 to 8 days.

Last Updated February 2022