

	Do not use in female dairy cattle 20 months of age or older. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.			
	Noromectin® Injection for Cattle and Swine ( <i>Norbrook</i> ) <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.	Cattle	S.C	35d
	<b>Noromectin® Plus Injection for Cattle</b> ( <i>AgriLabs</i> ) <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.	Cattle	S.Q.	49d
	Noromectin® Pour-On for Cattle ( <i>Norbrook</i> ) <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	Topical	48d
	<b>Noromycin 300 LA</b> ( <i>Norbrook</i> ) <b>Note:</b> Not for use in lactating dairy animals.	Cattle	I.M., I.V., S.Q.	28d
Prescription Required	<b>NuflorGOLD™</b> ( <i>Intervet/Merck Animal Health</i> ) <b>Note:</b> Do not use in calves to be processed for veal. Not for use in female dairy cattle 20 months of age or older.	Cattle	S.Q.	44d
Prescription Required	<b>Nuflor® Injectable Solution 300mg/mL</b> ( <i>Intervet/Merck Animal Health</i> ) <b>Note:</b> Do not use in calves to be processed for veal. Not for use in female dairy cattle 20 months of age or older.	Cattle Cattle	I.M. S.Q.	28d 38d
	<b>Nuplura™ PH</b> ( <i>Novartis (Farm Animal)</i> )	Cattle	S.Q.	60d
	<b>Ocu-guard® MB-1</b> ( <i>Beohringer Ingelheim</i> )	Cattle	S.Q.	21d
	<b>Once PMH® SQ</b> ( <i>Intervet/Merck Animal Health</i> )	Cattle	S.Q.	21d
	<b>One Shot®</b> ( <i>Zoetis</i> )	Cattle	I.M., S.Q.	21d
	<b>One Shot Ultra™ 7</b> ( <i>Zoetis</i> )	Cattle	S.Q.	21d
	<b>One Shot Ultra™ 8</b> ( <i>Zoetis</i> )	Cattle	S.Q.	21d
Prescription Required	<b>Orbenin-DC®</b> ( <i>Intervet/Merck Animal Health</i> ) <b>Note:</b> Not to be used within 28 days of calving.	Dairy Cattle	I.M.M.	28d
	<b>Orbeseal®</b> ( <i>Zoetis</i> )	Dairy Cattle	I.M.M.	0d 0h
	<b>Organic Iodide Powder</b> ( <i>Iowa Veterinary Supply Co.</i> )	Cattle	In the feed	

	<b>Note:</b> Do not administer to lactating dairy cattle.			
	<b>Oxytet 100 (Norbrook)</b> <b>Note:</b> Do not use in calves to be processed for veal. Not for use in lactating dairy animals.	Cattle	I.V.	22d
	<b>Oxytetracycline Injection 200 (Norbrook)</b>	Beef Cattle Dairy Cattle	I.M., I.V., S.Q. I.M., I.V., S.Q.	28d 28d 96h
	<b>Oxytocin (Generic)</b> <b>Note:</b> Not for human use. Women of childbearing age should handle with extreme caution.	Beef Cattle Dairy Cattle	I.M. I.M.	0d 0d
Prescription Required	<b>Panacur® Cattle Dewormer Suspension 10% (Intervet/Merck Animal Health)</b> <b>Note:</b> Do not use in calves to be processed for veal. Do not use at 10 mg/kg in dairy cattle. Dose rate of 10 mg/kg is for beef cattle only. Dose rate of 10 mg/kg in dairy cattle could result in violate residue in milk.	Cattle	Oral	8d
	<b>Pennox 343® Soluble Powder (PennField)</b> <b>Note:</b> A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.	Cattle	dw	5d
	<b>Piliguard® Pinkeye-1 Trivalent (Intervet/Merck Animal Health)</b>	Cattle	I.M., S.Q.	60d
	<b>Piliguard® Pinkeye Triview® (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	60d
	<b>Pinkeye Shield® XT4 (Novartis (Farm Animal))</b>	Cattle	I.M.	60d
Prescription Required	<b>Pirsue® Sterile Solution (Zoetis)</b>	Dairy Cattle	I.M.M.	9d 36h
Prescription Required	<b>Polyflex® (Boehringer Ingelheim)</b>	Beef Cattle Dairy Cattle	I.M. I.M.	6d 6d 48h
	<b>Polyotic® Soluble Powder (Boehringer Ingelheim)</b> <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	dw	4d
Prescription Required	<b>Predef® 2X Sterile Aqueous Suspension (Zoetis)</b> <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	I.M.	7d
	<b>PregGuard® GOLD FP® 10 (Zoetis)</b>	Cattle	I.M.	21d
	<b>Presponse® HM (Boehringer Ingelheim)</b>	Cattle	I.M.	21d

<b>Presponse® SQ (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Pro-Bac-C (AgriLabs)</b> <b>Note:</b> Do not use in calves to be processed for veal. Do not use with other sources of lasalocid.	Cattle	mr		
<b>Pro-Pen-G® Injection (Bimeda)</b>	Beef Cattle Dairy Cattle	I.M. I.M.	4-7d 4-7d	48h
<b>Prozap® Zipcide® Dust (Chem-Tech)</b>	Beef Cattle Dairy Cattle	Topical Topical	0d 0d	0h
<b>Pulmo-Guard™ MpB (Boehringer Ingelheim)</b>	Beef Cattle	S.Q.	60d	
<b>Pulmo-Guard® PH-M (AgriLabs)</b>	Cattle	S.Q.	21d	
<b>Pulmo-Guard™ PHM-1 (AgriLabs)</b>	Cattle	S.Q.	60d	
<b>Pyramid® 4 + Presponse® SQ (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Pyramid® 5 (Boehringer Ingelheim)</b>	Cattle	I.M., S.Q.	21d	
<b>Pyramid® 5 + Presponse® SQ (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Pyramid® 10 (Boehringer Ingelheim)</b>	Cattle	I.M., S.Q.	21d	
<b>Prescription Quartermaster® (Zoetis)</b> <b>Required Note:</b> Animals infused with this product must not be slaughtered for use in food within 96 hours after calving. Not for use in lactating cows. This product must not be used within 6 weeks of the expected date of calving.	Dairy Cattle	I.M.M.	60d	96h
<b>Quatracon-2X™ (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Ralgro® Implants for Beef Cattle (Intervet/Merck Animal Health)</b> <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not use in bulls intended for reproduction or dairy animals. Do not use before 1 month of age or after weaning in heifers intended for reproduction.	Beef Cattle	Implant		
<b>Repel-X® Insecticide &amp; Repellent (Farnam)</b> <b>Note:</b> Do not apply directly to any animals destined for milking.	Cattle	Premise		
<b>ReproSTAR® L5 HB (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d	

<p>Prescription Required <b>Resflor Gold®</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not use in female dairy cattle 20 months of age or older.</p>	Cattle	S.Q.	38d
<p><b>RESPISHIELD™ HM</b> (<i>Merial</i>)  <b>Resvac® 4/Somubac®</b> (<i>Zoetis</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not use in female dairy cattle 20 months of age or older.</p>	Cattle	S.Q.	60d
<p><b>Revalor®-200</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.</p>	Beef Cattle	Implant	
<p><b>Revalor®-G</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.</p>	Beef Cattle	Implant	
<p><b>Revalor®-H</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.</p>	Beef Cattle	Implant	
<p><b>Revalor®-IH</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.</p>	Beef Cattle	Implant	
<p><b>Revalor®-IS</b> (<i>Intervet/Merck Animal Health</i>)  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for</p>	Beef Cattle	Implant	

veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.

**Revalor®-S (Intervet/Merck Animal Health)**

**Note:** A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.

Beef  
Cattle

Implant

**Revalor®-XS (Intervet/Merck Animal Health)**

**Note:** A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food. Not to be used in dairy animals.

Beef  
Cattle

Implant

**Rumatel® 88 (Phibro)**

Beef  
Cattle  
Goats

In the feed  
In the feed

14d  
30d

0h

**Saber™ Extra Insecticide Ear Tags (Intervet/Merck Animal Health)**

**Note:** Remove tags before slaughter.

Cattle

Ear Tags

**Saber™ Pour-On Insecticide (Intervet/Merck Animal Health)**

**Note:** Do not apply to lactating or dry dairy cows. Not recommended for use on veal calves.

Beef  
Cattle

Topical

**Safe-Guard® Cattle Dewormer Paste 10% (Intervet/Merck Animal Health)**

Beef  
Cattle  
Dairy  
Cattle

Oral  
Oral

8d  
8d

0h

**Safe-Guard® Dewormer for Beef & Dairy Cattle and Goats Suspension 10% (Intervet/Merck Animal Health)**

**Note:** A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Beef  
Cattle  
Dairy  
Cattle

Oral  
Oral

8d  
8d

0h

**Safe-Guard® Medicated Dewormer for Beef & Dairy Cattle 0.5% Alfalfa-Based Pellets (Intervet/Merck Animal Health)**

Beef  
Cattle  
Dairy  
Cattle

In the feed  
In the feed

13d  
13d

0h

**Safe-Guard® Medicated Dewormer for Beef & Dairy Cattle, & Swine (0.5% Top Dress Pellets 10 lbs) (Intervet/Merck Animal Health)**

Beef  
Cattle  
Dairy

In the feed  
In the feed

13d  
13d

0h

	Cattle			
<b>Safe-Guard® Medicated Dewormer for Beef &amp; Dairy Cattle (Flaked Meal)</b> ( <i>Intervet/Merck Animal Health</i> )	Beef	In the feed	13d	0h
	Cattle	In the feed	13d	
	Dairy Cattle			
<b>Safe-Guard® Medicated Dewormer for Beef Cattle (20% Protein Block)</b> ( <i>Intervet/Merck Animal Health</i> )	Beef	Oral	16d	
	Cattle			
<b>Safe-Guard® Medicated Dewormer for Beef Cattle (En-Pro-Al® Molasses Block)</b> ( <i>Intervet/Merck Animal Health</i> )	Beef	Oral	11d	
	Cattle			
<b>Salmonella Newport Bacterial Extract</b> ( <i>Zoetis</i> )	Cattle	S.Q.	60d	
<b>Salmo Shield® TD</b> ( <i>Novartis(Farm Animal)</i> )	Cattle	I.M., S.Q.	21d	
<b>Scour Bos® 4</b> ( <i>Novartis (Farm Animal)</i> )	Cattle	I.M.	60d	
<b>Scour Bos® 9</b> ( <i>Novartis (Farm Animal)</i> )	Cattle	I.M.	60d	
<b>ScourGuard® 4 KC</b> ( <i>Zoetis</i> )	Cattle	I.M.	21d	
<b>Sodium Iodide 20%</b> ( <i>Generic</i> ) <b>Note:</b> Not for use in pregnant, lactating dairy cattle or animals that are exhibiting hypothyroidism.	Cattle	I.V.		
<b>SolidBac® Pinkeye IR/PR</b> ( <i>Zoetis</i> )	Cattle	S.Q.	21d	
<b>Somubac®</b> ( <i>Zoetis</i> )	Cattle	S.Q.	21d	
Prescription Required <b>Spectramast® DC</b> ( <i>Zoetis</i> )	Dairy Cattle	I.M.M.	0-16d	0h
Prescription Required <b>Spectramast® LC</b> ( <i>Zoetis</i> )	Dairy Cattle	I.M.M.	2d	72h
<b>Spirovac®</b> ( <i>Zoetis</i> )	Cattle	S.Q.	21d	
<b>Spirovac® L5</b> ( <i>Zoetis</i> )	Cattle	I.M., S.Q.	21d	
<b>Spirovac® VL5</b> ( <i>Zoetis</i> )	Cattle	I.M., S.Q.	21d	
<b>StandGuard® Premise Insecticide</b> ( <i>Elanco</i> ) <b>Note:</b> Do not apply to lactating or dry dairy cows. Not recommended for use on veal calves.	Cattle	Premise		
<b>Streptococcus Uberis Bacterin</b> ( <i>Hygieia</i> )	Dairy Cattle	S.Q.	60d	
<b>Sulfadimethoxine Injection - 40%</b> ( <i>Aspen</i> ) <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.	Beef	I.V.	5d	60h
	Cattle	I.V.	5d	
	Dairy			
	Cattle			
<b>Sulfadimethoxine Oral Solution</b> ( <i>Generic</i> ) <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	dw	7d	

<p><b>Sulfadimethoxine Soluble Powder (Generic)</b>  <b>Note:</b> A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.</p>	Beef Cattle	dw	7d
<p><b>Sulmet® Drinking Water Solution 12.5% (Boehringer Ingelheim)</b>  <b>Note:</b> Not for use in female dairy cattle 20 months of age or older.</p>	Cattle	dw	10d
<p><b>Sulmet® Oblets® (Boehringer Ingelheim)</b>  <b>Note:</b> Not for use in female dairy cattle 20 months of age or older.</p>	Cattle	Oral	10d
<p><b>Sulmet® Soluble Powder (Boehringer Ingelheim)</b>  <b>Note:</b> Not for use in female dairy cattle 20 months of age or older.</p>	Cattle	dw	10d
<p><b>Sustain III® (Generic)</b>  <b>Note:</b> Do not use in lactating dairy cattle. Not for use in female dairy cattle 20 months of age or older.</p>	Cattle	Oral	12d
<p><b>Sustain III® Calf Bolus (Generic)</b>  <b>Note:</b> Not for use in female dairy cattle 20 months of age or older.</p>	Cattle	Oral	12d
<p><b>Synanthic® Bovine Dewormer Suspension, 9.06% (Boehringer Ingelheim)</b></p>	Cattle	Oral	7d
<p><b>Synovex® C (Zoetis)</b>  <b>Note:</b> Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food.</p>	Beef Cattle	Implant	
<p><b>Synovex® Choice (Zoetis)</b>  <b>Note:</b> Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food.</p>	Beef Cattle	Implant	
<p><b>Synovex® H (Zoetis)</b>  <b>Note:</b> Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food.</p>	Beef Cattle	Implant	
<p><b>Synovex® Plus™ (Zoetis)</b>  <b>Note:</b> Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food.</p>	Beef Cattle	Implant	

<b>Synovex® S (Zoetis)</b> <b>Note:</b> Do not use in calves to be processed for veal. Implant in the ear only. Do not attempt to salvage the implanted ear for human or animal food.	Beef Cattle	Implant		
<b>Terramycin® Scours Tablets (Zoetis)</b> <b>Note:</b> Do not use in calves to be processed for veal. Not for use in lactating dairy animals.	Cattle	Oral	7d	
<b>Tetanus Antitoxin, Equine Origin (Colorado Serum)</b>	Cattle	I.M., S.Q.	21d	
<b>Tetracycline Soluble Powder 324 (AgriPharm)</b> <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	dw	5d	
<b>TetraMed® 324 HCA (Bimeda)</b> <b>Note:</b> Do not use in calves to be processed for veal.	Cattle	dw	5d	
<b>Titanium® 5 (AgriLabs)</b>	Cattle	I.M., S.Q.	21d	
<b>Titanium® 5 L5 HB (AgriLabs)</b>	Cattle	I.M., S.Q.	21d	
<b>ToDAY® (Boehringer Ingelheim)</b>	Dairy Cattle	I.M.M.	4d	96h
<b>ToMORROW (Boehringer Ingelheim)</b> <b>Note:</b> Milk from treated cows must not be used for food during the first 72 hours after calving. Not to be used within 30 days of calving.	Dairy Cattle	I.M.M.	42d	72h
<b>Triangle® 4 + HS (Boehringer Ingelheim)</b>	Cattle	I.M., S.Q.	21d	
<b>Triangle® 4 + PH/HS (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Triangle® 4 + PH-K (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Triangle® 5 (Boehringer Ingelheim)</b>	Cattle	I.M., S.Q.	21d	
<b>Triangle® 9 + HS (Boehringer Ingelheim)</b>	Cattle	I.M.	21d	
<b>Triangle® 9 + PH-K (Boehringer Ingelheim)</b>	Cattle	S.Q.	21d	
<b>Triangle® 10 (Boehringer Ingelheim)</b>	Cattle	I.M., S.Q.	21d	
<b>TSV-2® (Zoetis)</b>	Cattle	I.N.	21d	
<b>Tylan® 50 Injection (Elanco)</b> <b>Note:</b> Do not use in calves to be processed for veal. Do not use in dry dairy cows. Not for use in female dairy cattle 20 months of age or older.	Cattle	I.M.	21d	
<b>Tylan® 200 Injection (Elanco)</b> <b>Note:</b> Do not use in calves to be processed for veal. Do not use in dry dairy cows. Not for use in female dairy cattle 20 months of age or older.	Cattle	I.M.	21d	
<b>Ultrabac® 7 (Zoetis)</b>	Cattle	S.Q.	21d	



<b>Ultrabac® 7/Somubac® (Zoetis)</b>	Cattle	S.Q.	21d
<b>Ultrabac® 8 (Zoetis)</b>	Cattle	S.Q.	21d
<b>Ultrabac® CD (Zoetis)</b>	Cattle	S.Q.	21d
<b>UltraChoice® 7 (Zoetis)</b>	Cattle	S.Q.	21d
<b>UltraChoice® 8 (Zoetis)</b>	Cattle	S.Q.	21d
<b>Ultra Saber™ Pour-On Insecticide (Intervet/Merck Animal Health)</b>	Beef Cattle	Topical	
<b>Note: Do not apply to lactating or dry dairy cows. Not recommended for use on veal calves.</b>			
<b>Valbazen® Suspension (Zoetis)</b>	Cattle	Oral	27d
<b>Vib Shield® Plus L5 (Novartis (Farm Animal))</b>	Cattle	I.M.	60d
<b>Vira Shield® 6 (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d
<b>Vira Shield® 6+L5 (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d
<b>Vira Shield® 6+L5 HB (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d
<b>Vira Shield® 6+L5 HB Somnus (Novartis (Farm Animal))</b>	Cattle	I.M.	60d
<b>Vira Shield® 6+L5 Somnus (Novartis (Farm Animal))</b>	Cattle	I.M.	60d
<b>Vira Shield® 6+Somnus (Novartis (Farm Animal))</b>	Cattle	I.M.	60d
<b>Vira Shield® 6+VL5 (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d
<b>Vira Shield® 6+VL5 HB (Novartis (Farm Animal))</b>	Cattle	S.Q.	60d
<b>Vision® 7 Somnus with Spur® (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vision® 7 with Spur® (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vision® 8 Somnus with Spur® (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vision® 8 with Spur® (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vista® 5 L5 SQ (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vista® 5 SQ (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vista® 5 VL5 SQ (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vista® Once SQ (Intervet/Merck Animal Health)</b>	Cattle	S.Q.	21d
<b>Vita Charge® 28 Dispersible Powder (BioZyme)</b>	Cattle	In the feed, mr	0d
<b>Vita Charge® Bolus (BioZyme)</b>	Cattle	Oral	0d
<b>Prescription Vitamin AD Injection (Clipper (Phoenix Pharm.)) Required</b>	Cattle	I.M., S.Q.	60d

<b>Wart Vaccine (Colorado Serum)</b>	Cattle	S.Q.	21d	
<b>Y-TeX® Co-Ral® Livestock Dust (Y-TeX)</b>	Beef Cattle Dairy Cattle	Topical Topical	0d 0d	0h
<b>Y-TeX® GardStar® Plus Insecticide Ear Tags (Y-TeX)</b> <b>Note:</b> Remove tags before slaughter.	Cattle	Ear Tags		
<b>Y-TeX® PYthon® Insecticide Cattle Ear Tags (Y-TeX)</b> <b>Note:</b> Remove tags before slaughter.	Cattle	Ear Tags	0d	
<b>Y-TeX® PYthon® MagnuM™ Insecticide Cattle Ear Tags (Y-TeX)</b> <b>Note:</b> Remove tags before slaughter.	Beef Cattle Dairy Cattle	Ear Tags Ear Tags	0d 0d	0h
<b>Y-TeX® Warrior™ Insecticide Cattle Ear Tags (Y-TeX)</b> <b>Note:</b> Remove tags before slaughter.	Cattle	Ear Tags	0d	
<b>Prescription ZACTRAN® (Merial)</b> <b>Required Note:</b> Do not use in calves to be processed for veal. Do not use in female dairy cattle 20 months of age or older.	Cattle	S.Q.	35d	
<b>Prescription Zuprevo™ 18% (Intervet/Merck Animal Health)</b> <b>Required Note:</b> Do not use in calves to be processed for veal. Do not use in female dairy cattle 20 months of age or older.	Cattle	S.Q.	21d	

Last Updated JANUARY 2014