




# Cattle Dewormer Comparison

Product	Active Ingredient	Persistent Activity	Lungworms	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Lice	Mange Mites	Horn Flies	Grubs	Beef	Lactating Dairy	Non-Lactating Dairy	Calves	Comments	Dosage	Slaughter Withdrawal	Administration
<b>CYDECTIN Injectable</b>	Moxidectin	14-42 days	X								X			X	X		X	X	X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	21 days	SubQ
<b>CYDECTIN Pour-On</b>	Moxidectin	28-42 days	X								X			X	X	X	X	X	X	X	X	Do not use in calves to be processed for veal.	5 ml / 110 lbs	0 days	Topical
<b>DECTOMAX Injectable</b>	Doramectin	21-28 days	X								X	X	X	X	X		X		X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 100 lbs	35 days	SubQ/IM
<b>DECTOMAX Pour-On</b>	Doramectin	21-35 days	X	X	X		X		X	X	X	X	X	X	X	X	X	X	X	X	X	For use in dairy replacement heifers under 20 months of age. Not for use in veal calves.	50 ml / 1100 lbs	45 days	Topical
<b>Eprinex Pour-On</b>	Eprinomectin	21-28 days	X								X			X	X	X	X	X	X	X	X	Do not use in calves to be processed for veal.	1 ml / 110 lbs	0 days	Topical
<b>EpriGard</b>	Eprinomectin	21-28 days	X								X			X	X	X	X	X	X	X	X	Do not use in calves to be processed for veal.	1 ml / 110 lbs	0 days	Topical
<b>Eprizero</b>	Eprinomectin	21-28 days	X								X			X	X	X	X	X	X	X	X	Do not use in calves to be processed for veal.	1 ml / 110 lbs	0 days	Topical
<b>Ivermectin Pour-On</b>	Ivermectin	14-28 days	X								X			X	X	X	X	X	X	X	X	Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 22 lbs	48 days	Topical
<b>Ivermectin Injectable</b>	Ivermectin	14-28 days	X								X			X	X		X		X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	35 days	SubQ
<b>Ivomec Plus Injectable</b>	Ivermectin/ Clorsulon	14-28 days	X					X		X				X	X		X		X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	49 days	SubQ
<b>Ivomec Pour-On</b>	Ivermectin	14-28 days	X								X			X	X	X		X		X		Do not use in female dairy cattle over 20 months of age or veal calves.	50 ml / 1100 lbs	48 days	Topical
<b>Ivermectin Plus Injectable</b>	Ivermectin/ Clorsulon	14-28 days	X					X		X				X	X		X		X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	49 days	SubQ
 <b>LongRange Injectable</b>	Eprinomectin	100-150 days	X	X	X	X	X							X			X		X			Do not use in female dairy cattle over 20 months of age or veal calves.	1 ml / 110 lbs	48 days	SubQ
 <b>panacur 10%</b>	Fenbendazole		X	X	X	X	X											X	X	X	X	Do not use in calves to be processed for veal.	2.3 ml / 100 lbs*	8 days	Drench

# Cattle Dewormer Comparison

Product	Active Ingredient	Persistent Activity	Lungworms	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Lice	Mange Mites	Horn Flies	Grubs	Beef Lactating Dairy	Non-Lactating Dairy Calves	Comments	Dosage	Slaughter Withdrawal	Administration		
<b>PROHIBIT</b>	Levamisole		X	X	X	X	X												X	X	X	Do not use in female dairy cattle over 20 months of age.	2 ml / 100 lbs	2 days	Drench
<b>Prozap Backrubber and Pour-On</b>	Permethrin													X		X			X	X	X	Not recommended for use on veal calves.	1 oz / 100 lbs (max 8 oz)	0 days	Topical
<b>Prozap Protectus Pour On</b>	Diflubenzuron, Piperonyl butoxide, Lambda-cyhalothrin													X		X			X				3 ml / 100 lbs (max 30 ml)		Topical
<b>Rumatel Crumbles</b>	Morantel Tartrate		X							X									X	X	X		0.1 lb / 100 lbs	14 days	Top-dress
<b>safe-guard 0.5% Alfa-Based Pellets</b>	Fenbendazole		X	X	X	X	X												X	X	X		0.5 lb / 500 lbs	13 days	Top-dress
<b>safe-guard Medicated Dewormer (20% Protein)</b>	Fenbendazole		X	X	X	X	X												X	X	X	Do not use in female dairy cattle over 20 months of age.	1.5 lbs / 500 lbs	16 days	Block - Free Choice
<b>safe-guard Molasses Block</b>	Fenbendazole		X	X	X	X	X												X	X	X	Do not use in female dairy cattle over 20 months of age.	2.27 mg / lb	11 days	Block - Free Choice
<b>safe-guard Paste 10%</b>	Fenbendazole		X	X	X	X	X												X	X	X	Do not use in calves to be processed for veal.	5 g / 220 lbs	8 days	Oral
<b>safe-guard Suspension 10%</b>	Fenbendazole		X	X	X	X	X												X	X	X	Do not use in calves to be processed for veal.	2.3 ml / 100 lbs*	8 days	Drench
<b>safe-guard Type B Medicated (Flaked Meal)</b>	Fenbendazole		X	X	X	X	X												X	X	X		0.25 lb / 1000 lbs	13 days	Free Additive
<b>Synanthic Suspension</b>	Oxfendazole		X	X	X		X		X										X	X	X	Do not use in female dairy cattle over 20 months of age.	1 ml / 110 lbs	7 days	Drench
<b>Valbazen</b>	Albendazole		X	X	X	X	X	X	X	X									X	X	X	Do not use in female dairy cattle over 20 months of age.	4 ml / 100 lbs	27 days	Drench
 <b>Valcor Injectable</b>	Doramectin/ Levamisole	14-56 days	X	X	X	X	X				X		X	X	X		X		X			For use in beef cattle 2 months of age or older and in replacement dairy heifers less than 20 months of age.	1 ml / 55 lbs	15 days	SubQ

Persistent Activity - Proven to effectively protect cattle from re-infection of certain parasites.

\* 2.3 ml/100 lbs or 2.3 ml/50 lbs for ascarids

Last Updated October 2023